



ACC.15

TCT@ACC-12 | innovation in intervention

A1375
JACC March 17, 2015
Volume 65, Issue 10S

Prevention

ADDITION OF SPECIFIC LIPOPROTEIN(A) APHERESIS IN STABLE ISCHEMIC HEART DISEASE PATIENTS CONTRIBUTES TO CAROTID ATHEROSCLEROSIS PREVENTION

Poster Contributions

Poster Hall B1

Saturday, March 14, 2015, 10:00 a.m.-10:45 a.m.

Session Title: Lipids, Novel Therapies and Acute Coronary Syndromes

Abstract Category: 21. Prevention: Clinical

Presentation Number: 1107-110

Authors: *Marat V. Ezhov, Maya Safarova, Olga Afanasieva, Olga Pogorelova, Tatyana Balakhonova, Valery Kukharchuk, Sergei Pokrovsky, Russian Cardiology Research and Production Center, Moscow, Russian Federation*

Background: The purpose of this study was to evaluate the effect of 18-month course of specific lipoprotein(a) (Lp(a)) apheresis on carotid intima-media thickness (CIMT) progression in patients with controlled low-density lipoprotein cholesterol (LDL-C) levels reached with atorvastatin treatment, but with high Lp(a) levels, during 42 months of follow-up.

Methods: In this single-center, partially-blind study 30 stable ischemic heart disease patients with Lp(a) level ≥ 50 mg/dL on a guideline-driven therapy were assigned to Lp(a) immunoadsorption columns (n=15, Lp(a) Lipopak®, POCARD Ltd., Russia) on a weekly basis or atorvastatin monotherapy after a 4-week diet and atorvastatin run-in period. The primary efficacy end-point was the absolute change from baseline through 18 months in mean of left and right CIMT, as assessed by duplex ultrasound scanning.

Results: As the only target for the specific Lp(a) apheresis, Lp(a) level decreased by an average of $73 \pm 12\%$ (a mean for 951 procedures) to 29 ± 16 mg/dL immediately after the session. During 18 months of regular treatment, the interval mean values for Lp(a) were 73.3 ± 13.2 mg/dL. The final change in Lp(a) level in the immunoadsorption group was -31.7 ± 22.3 mg/dL, as compared with 4.8 ± 10.8 mg/dL in the atorvastatin only group ($P < 0.0001$). LDL-C level corrected for Lp(a) cholesterol unchanged: -7.7 ± 23.2 and -3.9 ± 23.2 mg/dL for apheresis and control groups, respectively ($P = 0.65$). At 18 months the primary efficacy end point decreased by -0.07 ± 0.15 mm ($P = 0.01$) in the Lp(a) apheresis group and by -0.03 ± 0.13 ($P = 0.09$) in the control group. Two years after apheresis termination, in the immunoadsorption-treated patients (n=14) Lp(a) level was 103 ± 23 mg/dL and CIMT increased in average by 0.02 ± 0.08 mm ($P = 0.12$ in comparison with the baseline). In the control group Lp(a) was 101 ± 52 mg/dL and CIMT increased by up to 0.06 ± 0.10 mm ($P = 0.18$ versus baseline, $P = 0.03$ for between-group comparison).

Conclusion: Specific Lp(a) apheresis during 18 months resulted in significant regression of CIMT in comparison with the traditional approach despite achieved LDL-C levels in both groups. Lp(a) lowering strategy led to long-term stabilization in the rate of CIMT progression.